



## Article

# Effect of Adding Some Substances in Media for Induction and Development of Callus

Hiba N. A. AlAkaidi

Anesthesia Technique Department, Medical Technical Institute of Mosul, Northern Technical University, Iraq.



CrossMark

\*Corresponding author: [hiba.nawaf@ntu.edu.iq](mailto:hiba.nawaf@ntu.edu.iq)

Future Science Association

Available online free at  
[www.futurejournals.org](http://www.futurejournals.org)

Print ISSN: 2692-5826

Online ISSN: 2692-5834

DOI:

10.37229/fsa.fjh.2024.09.17

Received: 27 July 2024

Accepted: 24 August 2024

Published: 17 September 2024

**Publisher's Note:** FA stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** Callus is a mass of unspecialized parenchyma cells, and its size and growth increase depending on the type and quantity of growth regulators present in the nutrient medium. This tissue is commonly used for plant propagation through tissue culture by inducing and growing it on media containing auxins, cytokinins, amino acids, sucrose, salicylic acid, and the effect of these additions on enhancing callus induction.

**Key words:** Callus, Salicylic acid, Amino acids, Sucrose, tryptophane.

## Introduction

Research studies have shown that the basic structure of alkaloids, particularly the tropane group, is derived from certain amino acids and other biological molecules (Al-Shahat, 2006). To increase the production of secondary metabolites, it is essential to add the intermediate compounds that produce them, including amino acids. It has been observed that their addition has increased the production of tropane alkaloids (Silvestrini *et al.*, 2002; Whitmer *et al.*, 1998; Moreno *et al.*, 1993).

## Tryptophan

Tryptophan is one of the important amino acids involved in the biosynthesis of alkaloids, which are primary metabolic products from which alkaloid groups are formed, providing the carbon skeleton and nitrogen components for alkaloids (Sarin, 2005; Al-Saadawi and Younis, 1991). This was confirmed by Dmitruk (1973) in his study on the effect of amino acids on the production of tropane alkaloids from the plant *Scopolia* sp. It was

found that the amino acid Tryptophan was the most effective in producing the alkaloid Hyoscyamine and the alkaloid Scopolamine.

### Salicylic Acid

The SA is embodied in plant growth and development, as it affects photosynthesis, transpiration, and the absorption and transport of nutrients within the plant (Sakhanokho and Kelley, 2009; Hayat and Ahmad, 2007). It also induces or causes distinctive changes in the anatomy of the plant leaf and the structure of chloroplasts. It plays an important role in inducing plant resistance against pathogens by stimulating plant cells to produce Pathogenesis-Related Proteins (Verberne *et al.*, 1999; Yalpani *et al.*, 1993). Furthermore, salicylic acid is involved in the acquired systemic resistance that occurs as a result of a pathogen attack in one part of the plant, and this signal can be transmitted to neighboring plants, prompting them to develop resistance by converting salicylic acid into the volatile ester methyl salicylate (Hayat and Ahmad, 2007; Molinari, 2007). Eskandari *et al.* (2012) studied the effect of adding SA at different concentrations on the alkaloids hyoscyamine and scopolamine from genetically modified hairy root cultures of the belladonna plant (*Atropa belladonna*) in MS medium, obtaining the highest amount of the mentioned alkaloids from the culture at 34 mg $\cdot$ L<sup>-1</sup> SA, which were 1.6 mg/gram hyoscyamine and 1.3 mg/gram scopolamine. Pitta *et al.* (2000) demonstrated that the use of SA at different concentrations led to an increase in the accumulation of hyoscyamine and scopolamine alkaloids from the hairy root culture of *Brugmansia candida* in MS medium. Al-Muhari (2014) managed to obtain the highest amount of callus, 0.584 g, from the leaf culture of the periwinkle plant (*Catharanthus roseus*) in MS medium supplemented with 0.75 mg $\cdot$ L SA and 0.75 mg $\cdot$ L 2,4-D after 50 days of culture.

### Sodium chloride

Sodium chloride is a salt that increases the negative osmotic potential of the cell, thereby enhancing the exchange of solutes within the cell, which plays a role in increasing the process of photosynthesis and consequently the production of secondary metabolites. The composition of the nutrient medium supplemented with sodium chloride has profound effects on cell growth and the accumulation of secondary metabolic products, although many of these products are not associated with cell growth. The term "Elicitation" refers to the treatment of plant cells with biotic or abiotic stress factors, which is an effective means of improving the production of secondary metabolites in plant tissue culture. This strategy is based on the premise that this accumulation is part of the defensive reactions of plants against pathological infections and environmental stresses (Zhou and Wu, 2006). Numerous experiments have demonstrated the effect of adding sodium chloride to the culture medium on biological processes. Aljibouri *et al.* (2012) indicated that the addition of 200 mg/L sodium chloride to the MS nutrient medium for the callus induced from the plant *Hyoscyamus niger* L. resulted in a significant increase in the amount of Scopolamine and a decrease in the Hyoscyamine compound compared to the control treatment. Abdeil Rahman *et al.* (2013) in a study on the production of Atropine and Scopolamine in the plants *Daturai metel* and *Daturai stramonium*

found that the addition of 7 mg/L sodium chloride to the MS nutrient medium led to an increase in the compounds Atropine and Scopolamine in the induced callus compared to the control treatment without sodium chloride after one month of cultivation. **Al-Muhari (2014)** indicated that the highest amount of callus, 0.659 g, was obtained from the cultivation of leaves of the plant *Catharanthus roseus* on the MS nutrient medium supplemented with NaCl.

### Sucrose

Sucrose is the most commonly used sugar as an energy source in the propagation of plants outside the living organism. It is a non-reducing ketonic disaccharide composed of one glucose unit and one fructose unit (**Al-Bahr et al., 1999**). **Laszol (2003)** found that the highest fresh weight of the callus was when the root hairs of the *Datura innoxia* plant were cultured on MS nutrient medium supplemented with 10 and 20 g/l sucrose. **Iranbakhsh et al. (2006)** indicated that the highest production of alkaloids from the *Datura stramonium* plant was from the callus induced on MS nutrient medium supplemented with 20 g / l sucrose after 5 weeks of cultivation. **Aljibouri et al. (2012)** showed that culturing the induced callus of the *Hyoscyamus niger* L. plant on MS medium supplemented with 50 g/l sucrose resulted in a significant increase in the amount of Scopolamine and a decrease in the amount of Hyoscyamine compared to the control treatment.

### References

- Abdel Rahman, R.; Gomma, S. E.; Abdel Salam, N.R.; El-Wakil, H.M.F.; Khaled, A.S. and Hassan, H.M. (2013)**. Effect of sodium chloride on tropan alkaloids accumulation and proline content in *Datura metel* and *D. stramonium* callus cultures. International Journal of Advanced Biological and Biomedical Research, 1(2): 197-210.
- Al-Bahr, M. K.; F.A.R. Ahmed and Saqr, M.A. (1999)**. Plant Biotechnology. Tissue Culture and Genetic Engineering. Arab Publishing and Distribution Company. Cairo - Egypt.
- Aledo, J. C. (2004)**. Glutamine breakdown in rapidly dividing cell: waste or investment. Bio. Essays, 26 (7): 136-141.
- Aljibouri, A. M. J.; Al- Samarra, K. W.; Abd, A. S.; Mageed, D. M. and Ali, A. J. A. (2012)**. Alkaloids production from callus of *Hyoscyamus niger* L. *In vitro*. Journal of Life Sciences, 6: 874 – 882.
- Al-Maamari, A. M. S. (2014)**. Induction and Differentiation of Callus from Some Parts of the Plant *Catharanthus roseus* L. and Estimation of Its Alkaloid Content. Doctoral Thesis, College of Agriculture and Forestry / University of Mosul - Iraq.
- Al-Saadawi, I. S. and Younis, M. A. (1991)**. Nitrogen Metabolism in Plants. Ministry of Higher Education and Scientific Research / University of Baghdad - Iraq.
- Al-Shahat, N. A. Z. (2006)**. Physiology and Chemistry of Alkaloids in Medicinal Plants and Their Pharmaceutical and Therapeutic Importance. Dar Al-Kutub Al-Ilmiyya for Publishing and Distribution, Cairo, Egypt.

- Brosnan, J. T. (2003).** Interorgan amino acid transport and its regulator. *J. Nutr.* 133 (6 Suppl 1): 2068S – 2072S. PMID 12771367. <http://jn.nutrition.org/cgi/content/full/133/6/2068S>.
- Dmitruk, S. E. (1973).** Role of amino acids in the formation of Tropane alkaloids, *Mter. Obl. Nauchin. Konf-Beca. Khim. O.Va., posuyashch 1975 - letiya Khim – technol Toeusk poliekh. Ins.* 3rd 1972 (pub. 1973), 57-59.
- Eskandari, S.A.; Piri, K. H.; Kayhanfar, M. and Hasanloo, T. (2012).** Influence of jasmonic acid, yeast extract and salicylic acid on growth and accumulation of Hyoscyamine and Scopolamine in hairy root cultures of *Atropa belladonna* L. *International Journal of Agriculture*, 2 (4): 403 – 409.
- Hall, J. E. and Guyton, C. A. (2006).** *Textbook of Medical Physiology* (11<sup>th</sup> ed.). St. Louis, Mo: Elsevier Saunders. P. 393.
- Hayat, S. and Ahmed, A. (2007).** *Salicylic Acid: A Plant Hormone*" Springer. Dordrecht, The Netherlands.
- Iranbakhsh, A.; Oshaghi, M. A. and Magd, A. (2006).** Distribution of atropine and Scopolamine in different organs and stages of development in *Datura stramonium* L. (Solanaceae) structure and ultrastructure of biosynthesizing cells. *Acta Biologica Cracoviensia Series Botanica*, 48 (1): 13 -18.
- Laszol, I. (2003).** Investigation on primary and secondary metabolism of *Datura innoxia* Mill. tissue culture. Semmelweis University, Institute of Pharmacognosy, Budapest.
- Molinari, S. (2007).** New developments in understanding the role of salicylic acid in plant defense. *Nutrition and Natural Resources*, 2 (67): 10.
- Moreno, P. R. H.; Heijden, V. D. and verpoote, R. (1993).** Effect of terpenoids precursor feeding and elicitation on formation of indole alkaloids in cell suspension cultures of *Catharanthus roseus*. *Plant Cell Reports*, 12: 702 -705.
- Pitta, A.S. I.; Spollansky, T.C. and Giulietti, A. M. (2000).** The influence of different biotic and abiotic elicitors on production and profile of Tropane alkaloids in hairy root cultures of *Brugmansia candida* enzyme microb. *Technol.*, 26: 252-258.
- Sakhanokho, H. F. and Kelley, R. Y. (2009).** Influence of salicylic acid on *in vitro* propagation and salt tolerance in *Hibiscus acetosella* and *Hibiscus moscheutos* (Cv. 'Luna Red'). *African Journal of Biotechnology.*, 8(8): 1474-1481.
- Sarin, R. (2005).** Useful metabolites from plant tissue cultures. *Biotechnology*, 4 (2): 79-93.
- Silvestrni, A.; Pasqua, G.; Botta, B.; Monacelli, B.; Heijden, R.V. and Verpoote, R. (2002).** Effect of alkaloid precursor feeding on *acamptonheca acuminata* cell line. *Plant Physiology and Bio –Chemistry*, 40: 749-753.
- Taha, H.S.; El-Baher, M. K. and Seif - El-Naser, M. M. (2009).** *In vitro* studies on Egyptian *Catharanthus roseus* (L.) G. Don. IV: Manipulation of some amino acids as precursors for enhanced of indole alkaloids production in suspension cultures. *Aust. J. Basic and Appl. Sci.*, 3(4): 3137 – 3144.

**Verberne, M.C.; Budi Muljono, R.A. and Verpoet, R. (1999).** Salicylic Acid Biosynthesis. Biochemistry and Molecular Biology of Plant Hormones. Elsevier Science, B.V.(Eds).

**Whitmer, S.; Canel, C.; Hallard, D.; Goncalves, C. and Verpoote, R. (1998).** Influence of precursor availability on alkaloids accumulation by trans-genic cell line of *Catharanthus roseus*. Plant Physiology, 116: 853-857.

**Yalpani, N.; Leon, J.; Lawton, M.A. and Raskin, I. (1993).** Pathway of salicylic acid biosynthesis in healthy and virus – inoculated tobacco. Plant Physiol., 103: 315 – 321.

**Yuneva, M.; Zamboni, N.; Oefner, P.; Sachidanandam, R. and Lazebnik, Y. (2007).** Deficiency in glutamine but not glucose induces MYC- dependent apoptosis in human cells. J. of Cell Biology, 178 (1): 93-105.

**Zhou, L. G. and Wu, J. Y. (2006).** Development and application of herbal medicine in China. Nat. Prod. Rep., 23: 789-810.

### الخلاصة

يعد الكالس تجمع من خلايا برنكيمية غير متخصصة ويزداد حجم هذا الكالس ونموه وفق نوع منظمات النمو وكميتها الموجودة في الوسط الغذائي وعادة يستخدم هذا النسيج في إكثار النباتات بالزراعة النسيجية من خلال استحداثه وتنميته على الأوساط الحاوية على الاوكسينات والساييتوكاينينات والاحماض الأمينية والسكرور والكلوتامين وحامض الساليسليك وكلوريد الصوديوم ومدى تأثير هذه الإضافات في زيادة استحداث الكالس.